



Childhood Cancer Profile

South America

Burden of Childhood Cancer



63.6% of all childhood

cancer cases in

the LAC region



61.5%

of all childhood cancer deaths in the LAC region



8.6%

of all deaths among children and adolescents (1-19 years) are due to Cancer 18,475 new cases/year



cancer deaths among people <20 years, each year

Annual trends in childhood cancer mortality (2000-2012)





+0.16%



+0.19%

Proportion of Prioritized Cancers

ALL Burkitt Hodgkin (26.3%) (1.7%) (3.5%)

(3.9%)

RB



Wilms



LGG

(6.9%)

Cancer Registration and Survival



67

2

necialized

specialized pediatric PBCR

24.2%

of the population covered by populationbased cancer registries (PBCR) Mortality/
Incidence Ratio

For every 100 children diagnosed with cancer

33 die of the disease each year

Net Survival

All cancers

60%

Acute Lymphoblastic Leukemia

30-89%

Health System Response



72%

of the WHO essential medicines to treat children with cancer are included in the EML of the countries in the region



1 center/554,000 children and adolescents

236

centers to deliver specialized care for pediatric cancer



76

Universal Health Care Index for Service Coverage



60% / 80%

Percentage of countries with Early Detection Programme or Guideline/ Defined Referral System

Palliative Care

107 Pediatric Palliative Care
Teams



1.1 Teams/million of children <15 years