

Pan American Health

World Health Organization

Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory with Autochthonous Transmission

2013-2014 (to week noted)

Cumulative cases

Epidemiological Week / EW 23 (Updated 06 June 2014)

Country/Territory	Week ^a	Cases ^b		Incidence Rate ^c	Deaths	Population ^d
		Suspected	Confirmed	incluence Nate	Deaths	X 1000
Latin Caribbean						
Cuba					0	11,266
Dominican Republic ^{&}	Week 22	52,976	17	509.4	1	10,404
French Guiana [*]	Week 22	0	246	98.8	0	249
Guadaloupe	Week 22	28,320	1,328	6,362.2	1	466
Haiti [*]	Week 23	11,802	14	114.5	0	10,317
Martinique	Week 22	31,720	1,515	8,226.5	9	404
Puerto Rico	Week 22	0	1	0.0	0	3,688
Saint Barthelemy	Week 22	540	135	7,584.3	0	9
Saint Martin (French part)	Week 22	3,340	793	11,580.3	3	36
Subtotal		128,698	4,049	360.3	14	36,839
Non-Latin Caribbean						
Anguilla	Week 13		33	206.3	0	16
Antigua and Barbuda	Week 20		4	4.4	0	90
Aruba			0	0.0	0	109
Bahamas			0	0.0	0	377
Barbados			0	0.0	0	289
Cayman Islands			0	0.0	0	54
Curacao			0	0.0	0	147
Dominica	Week 20	1,817	122	2,656.2	0	73
Grenada			0	0.0	0	110
Guyana*	Week 23		16	2.0	0	800
Jamaica			0	0.0	0	2,784
Montserrat			0	0.0	0	5
Saint Kitts and Nevis	Week 21		22	43.1	0	51
Saint Lucia [^]	Week 23	214	30	149.7	0	163
Saint Vincent and the Grenadines	Week 23	212	67	270.9	0	103
Sint Maarten (Dutch part) [≡]	Week 12		123	307.5	0	40
Suriname			0	0.0	0	539
Trinidad and Tobago			0	0.0	0	1,341
Turks and Caicos Islands			0	0.0	0	48
Virgin Islands (UK)	Week 19	0	20	62.5	0	32
Virgin Islands (US)			0	0.0	0	105
Subtotal		2,243	437	36.8	0	7,276
TOTAL		130,941	4,486	307.0	14	44,115

NOTES

^a Epidemiological Week for which information is available.

^b **Suspected case**: patient with acute onset of fever >38°C (101°F) and severe arthralgia or arthritis not explained by other medical conditions, and who resides or has visited epidemic or endemic areas within two weeks prior the onset of the symptoms. **Confirmed case**: a suspected case with any specific CHIK test (viral isolation, RT-PCR, Ig M, or four-fold increase of CHIK specific antibodies titers) - PAHO/CDC confirmed case definition available at

^c Incidence / 100,000 pop.

^d PAHO/WHO. Basic Indicators, 2013. http://www.paho.org/hq/index.php?option=com_docman&task=doc_download&gid=23535<emid=270&lang=en , except St. Barthelemy ("Populations légales 2009 entrent en vigueur le 1er janvier 2012." Recensement de la population. INSEE. Retrieved 2 January 2014). Data source : Cases reported by IHR NFPs to PAHO/WHO and/or through Member States websites or official news publication

* As published by DIGEPI http://digepisalud.gob.do/boletines/boletines-semanales/doc_download/356-.html

* Includes 86 (35%) imported cases.

** Information on confirmed cases provided by CARPHA (6 Jun 2014).

Includes one imported case (EW 06 2014).

Data published by Le point épidémiologique, N°11, 2014.

[^]Includes one imported case (EW 12 2014). Additional four cases confirmed by CARPHA (EW 19 2014).

Pan American Health Organization World Health Organization REGOLAL OFFICE FORTHE Americas	Number of Imported Cases of Chikungunya Fever in the Americas, by Country or Territory Weekly and Cumulative cases Epidemiological Week / EW 23 (Updated 06 June 2014)				
Country/Territory	Week ^a	Weekly Imported Cases	Cumulative Imported Cases		
North America					
United States of America	Week 22		27		
Central America Isthmus	•				
Panama	Week 21	2	2		
Latin Caribbean					
Cuba	Week 21	2	2		
Southern Cone					
Chile	Week 22	1	2		
Non-Latin Caribbean					
Aruba	Week 06	1	1		
Barbados	Week 23	1	1		
TOTAL		7	35		

NOTES

^a Epidemiological Week for which information is available.

* Only imported cases in countries or territories **without** authoctonous trasnmsision; full epidemiology report on "Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory with Autochthonous **Data source** : Cases reported by IHR NFPs to PAHO/WHO and/or through Member States websites or official news publication