


COLOMBIA



RISK FACTORS			
Prevalence of:	Total (%)	Male (%)	Female (%)
Current tobacco smokers among adults ^{2,a}	18.0	31.0	5.0
Current use of any tobacco product among youth ²	27.6	27.0	27.8
Fruit and vegetable intake in adults ^{1,b}	•••		
Alcohol consumers, past 12 months ³	80.6	86.3	75.3
Low physical activity in adults ^{1,c}	79		
Obesity among adults (BMI ≥30) ¹	12.8	19.5	7.4
HPV prevalence (women with normal cytology) ⁴	•••		15.9

Notes: a: age-standardized prevalence of tobacco smoking in adults (15 years+), 2011; b: % who eat 5 or more servings of fruit and vegetables per day; c: % who are physically active for <600 met minutes; BMI: Body Mass Index; HPV: Human Papillomavirus.





Cancer type	Sex	Number of deaths (2009)	ASMR (2009)	AAPC
All cancer types	Male	19,410	128.31	-0.31
	Female	21,878	113.0	-0.93*
Breast cancer		3,030	15.18	1.00
Cervical cancer		2,169	10.75	-2.56*
Colorectal cancer	Male	1,441	9.39	2.88*
	Female	1,809	9.52	1.23
Liver cancer	Male	890	6.01	-0.56
	Female	1,096	5.86	-1.82*
Lung cancer	Male	2,820	19.28	0.51
	Female	1,981	10.69	0.5*
Prostate cancer		2,684	20.30	1.56*
Stomach cancer	Male	3,287	21.76	-1.81*
	Female	2,275	11.92	-3.06*

Notes: ASMR: Age-standardized mortality rate; AAPC: Average annual percent change; *: The AAPC is significantly different from zero at p=0.05.

CANCER PLANS AND CANCE	ER REGISTRATION ⁶
National integrated NCD policy, strategy or action plan	
Is there a national integrated NCD policy, strategy or action plan?	Yes
Is cancer included in the overall NCD plan?	Yes
First year of implementation	2011
Cancer policy, strategy or action plan	
Is there a cancer policy, strategy or action plan?	Yes
First year of implementation	2013
Cancer registry	
Is there a cancer registry?	Yes
Scope (national vs. subnational)?	Both
Type of registry (population vs. hospital based)?	Hospital based
Last year with available data	2008

CANCER PRIMARY PREVEN	NTION POLICIES
Tobacco control ²	
Smoke free environments?	All public spaces are completely smoke free
Raised taxes on tobacco?	Taxes are 44% of the final price
Health warning labels about the dangers of tobacco?	Medium size warning with all appropriate features or large warning lacking one or more appropriate features
Bans on tobacco advertising, promotion and sponsorship?	Ban on all forms of direct and indirect advertising
Overweight and obesity prevention and control ⁶	
Is there a policy, strategy or action plan for reducing overweight/obesity? (First year of implementation)	Under development
Harmful use of alcohol prevention and control ⁶	
Is there a policy, strategy or action plan for reducing harmful use of alcohol? (First year of implementation)	Yes (2011)
HPV vaccination in national immunization program? ⁷	Yes
Year of introduction/target age group/delivery strategy	2012/4 th grade school girls aged 9 years or older*/schools*
Estimated 3-dose coverage (calendar year)	
HBV vaccination in national immunization program?8	
Year of introduction/target age group	1994/newborn-2m-4m-6m; risk groups
Estimated 3-dose coverage (2012)	92%

Note: *HPV vaccination is introduced in two phases: the first phase targets fourth grade school girls aged 9 years and older, and the second phase will include girls out of schools in rural and urban areas.

CANCER SCREENING		
Cervical cancer		
Guidelines ^{9,10}		
Are there cervical cancer screening guidelines available?	Yes	
Guideline recommendations (tests/age groups/frequency)	Pap test/21-69 years of age/every 3 years after 2 consecutive normal annual Pap tests HPV test/30-65 years of age/every 5 years VIA/30-50 years of age/every 3 years	
Availability and coverage of screening services		
Cervical cytology ⁶	Generally available in the public and the private sectors	
Visual inspection with acetic acid ⁶	Generally available in the public and the private sectors	
Screening coverage ¹⁰	79% women aged 25-69 with a Pap test in 2010	

CANCE	R SCREENING
Breast cancer	
Guidelines ¹¹	
Are there breast cancer screening guidelines available?	Yes
Guideline recommendations (tests/age groups/frequency)	CBE/women aged 30 years or older/Annual Mammography/every 2 years in women aged 40-50 years/annual in women aged 50 years and older
Availability and coverage of screening services	
Clinical breast examination ⁶	Generally available in the public and the private sectors
Mammography ⁶	Generally available in the public and the private sectors
Screening coverage ¹²	34.2% CBE (2009); 54% mammography (2009)*
Colorectal cancer	
Guidelines ¹³	
Are there colorectal cancer screening guidelines available?	Yes
Guideline recommendations (tests/age groups/frequency)	Fecal occult blood test/ 50 years and older/every 2 years
Availability and coverage of screening services	i.
Fecal occult blood test or fecal immunological test ⁶	Generally available in the public and the private sectors
Exam or colonoscopy ⁶	Generally available in the public and the private sectors
Screening coverage	

Note: * Coverage data for Bogota D.C. and neighboring municipalities, based on a telephone survey

CANCI	R TREATME	NT AND PALLIATIVE CARE		
Radiotherapy ¹⁴		Oral morphine ⁶		
Availability of services in the public health sector? 6	Generally available	Availability in the public health sector?	Generally available	
Total energy teletherapy units/million inhabitants	1.4	Covered by health insurance or publically funded?	Yes	
Total radiotherapy centers	44	Appears in the national list of essential medicines?	Yes	
Radiation oncologists	87	Opioid consumption ¹⁵		
Medical physicists	49	Total morphine equivalence, per capita	6.74	
Chemotherapy ⁶		Community/home care ⁶		
Availability of services in the public health sector?	Generally available	Are community/home care services available in the public sector for people with advanced stages of NCDs?	Generally available	

Notes: ...: Information not available; N/A: Not applicable; m: months; VIA: Visual inspection with acetic acid; CBE: Clinical breast examination.

Sources:

- (1) PAHO. Non communicable Diseases in the Americas: Basic Health Indicators 2011. Washington D.C, USA, 2011.
- (2) PAHO. Tobacco Control Report for the Region of the Americas, 2013. Washington D.C, USA, 2013.
- (3) WHO. Global Information System on Alcohol and Health, retrieved August 28,2013, from http://bit.ly/17zhuqZ.
- (4) ICO. Information Center on HPV and Cancer, retrieved August 28, 2013, from http://www.hpvcentre.net/.
- (5) PAHO/WHO Mortality Database, 2012.
- (6) PAHO/WHO Survey on National Capacity for the Prevention and Control of Non-communicable Diseases, 2013.
- (7) Pediatric Society of Colombia. Pediavoz. Boletín informativo de la Sociedad Colombiana de Pediatría. Julio 2012, retrieved August 28, 2013 from http://bit.ly/XDp06U.
- (8) PAHO/WHO Comprehensive Family Immunization Unit. Immunization Country Profiles, retrieved August 28, 2013, from http://bit.ly/1dn4Td3: Immunization Reported Coverage in the Americas, by vaccine, retrieved August 28, 2013 from http://bit.ly/1dn4Td3: Immunization Reported Coverage in the Americas, by vaccine, retrieved August 28, 2013 from http://bit.ly/1dn4Td3: Immunization Reported Coverage in the Americas, by vaccine, retrieved August 28, 2013 from http://bit.ly/1dn4Td3: Immunization Country Profiles, retrieved August 28, 2013, from http://bit.ly/1dn4Td3: Immunization Reported Coverage in the Americas, by vaccine, retrieved August 28, 2013 from http://bit.ly/1dn4Td3: Immunization Reported Coverage in the Americas, by vaccine, retrieved August 28, 2013 from http://bit.ly/1dn59ZK.
- (9) Recomendaciones para la tamización de neoplasias del cuello uterino en mujeres sin antecedentes de patología cervical (preinvasora o invasora) en Colombia. Guías de práctica clínica (No.3). Available from: http://bit.ly/1cYxDv2.
- (10) Network of National Cancer Institutes of the South American Union (RINC/UNASUR). Informe preliminar reducido sobre la situación del control del cáncer de cuello uterino en 8 países de Latinoamérica. 2012
- (11) National Cancer Institute of Colombia. Recomendaciones para la tamización y la detección temprana del cáncer de mama en Colombia. Bogotá: INC, 2006, retrieved August 28, 2013 from http://bit.ly/12mNOLM.
- (12) Arboleda W et al. Cobertura de examen clínico y mamografía de tamización para cáncer de mama en mujeres bogotanas. RevColombCancerol 2009;13(2):69-76.
- (13) National Cancer Institute of Colombia. Guía de práctica clínica explícita basada en la evidencia de tamización de cáncer colorrectal para Colombia. 201, retrieved August 28,2013 from https://bit.ly/14TX7QI.
- (14) WHO/IAEA Directory of Radiotherapy Centers (DIRAC) Database, retrieved August 28, 2013 from http://www-naweb.iaea.org/nahu/dirac/.
- (15) University of Wisconsin. WHO Collaborating Centre for Pain Policy and Palliative Care, Opioid Consumption Database 2010, retrieved August 28, 2013, from http://www.painpolicy.wisc.edu/countryprofiles/amro.