

2010 Immunization Schedule for Selected Vaccines - Latin American Countries

Country	BCG		Polio						DPT				DPT-Hep B+Hib				DPT+Hib				Hep B**				Hib						
	1	2	NB§	1	2	3	4	5	6	1	2	3	4	5	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	
ARG*	NB			2m	4m	6m	18m	6y						6y	2m	4m	6m				18m	NB	11y (school children)								
BOL*	NB			2m	4m	6m									2m	4m	6m														
BRA*	NB			2m	4m	6m	15m						15m	4-6y					2m	4m	6m		NB	1m	6m						
CHI	NB			2m	4m	6m	18m						18m	4y	2m	4m	6m														
COL	NB			2m	4m	6m	18m	5y					18m	5y	2m	4m	6m					NB	risk groups								
COR	NB			2m	4m	6m	4y				4m	15m	4y		2m	6m						NB	risk groups; health workers				4m	15m			
CUB	NB			2 (<1y)	2 (1y)	2 (2y)	1 (9y)						18m	6y*	2m	4m	6m					NB									18m
DOR	NB			2m	4m	6m	18m	4y				18m	4y		2m	4m	6m					NB									
ECU	NB			2m	4m	6m	18m						18m	6y*	2m	4m	6m					NB*	11y (school children); health care workers								
ELS	NB			2m	4m	6m	12m	4y					12m	4y	2m	4m	6m					NB	risk groups								
GUT	NB			2m	4m	6m	18m	4y					18m	4y	2m	4m	6m					NB									
HAI	NB		NB	6w	10w	14w				6w	10w	14w																			
HON	NB			2m	4m	6m	18m						18m	4y	2m	4m	6m					NB	risk groups								
MEX*	NB			2m*	4m*	6m*	18m*							4y					2m*	4m*	6m*	18m*	NB	2m	6m	12y+					
NIC	NB			2m	4m	6m	*						18m	6y	2m	4m	6m						health workers								
PAN	NB			2m	4m	6m	18m	4y						4y	2m	4m	6m				18m	NB	10-11y; health workers at risk								
PAR	NB			2m	4m	6m	18m	4y					18m	4y	2m	4m	6m						risk groups								
PER	NB			2m	4m	6m							18m	4y	2m	4m	6m					NB	population at risk								
URU	NB			2m	4m	6m	12m							5y	2m	4m	6m	15m					12y	+1m	+6m	health workers					
VEN	NB			2m	4m	6m	18m	5y						5y	2m	4m	6m	18m					NB	risk groups							

text bold and in red are still pending replies

NB: Newborn

WCBA(s): Women of Childbearing Age

** This table only includes the hepatitis B vaccine used in children national schedules. Vaccination of at-risk groups is not included.

*** TT/dT use reported as indicated by countries in the EPI Indicator Tables.

§ Doses administered to newborns do not count towards the completion of the primary series.

* BOL: Hepatitis B for newborns only in selected hospitals.

* BRA: BCG at 6-10 years only in some states; Introduced rotavirus vaccine in 2006.

* CUB: DT

* ECU: HepB given to newborns in municipalities at risk; DT administered at 6 years

* MEX: Pentacel (DTaP-Hib-IPV) given; OPV doses for children 6 months to 5years old during National Health Weeks

* NIC: Additional OPV doses for children during National Health Weeks.

DISCLAIMER: Table not yet approved by countries

Updated: 28 September 2011



2010 Immunization Schedule for Selected Vaccines - Latin American Countries

Country	M	MMR			MR	TT/dT***					Yellow Fever†	Influenza		Varicella	Hepatitis A	
	Number of Doses															
	1	1	2	3	1	1	2	3	4	5	1	1	2	1	1	2
ARG*		1y	6y			16y	every 10y; pregnant women				≥1y in population in border areas	6m-23m with chronic diseases ≥65y		risk groups	12m	
BOL*		12-23m				WCBAs					12-23m					
BRA*		1y	4-6y			>7y	WCBAs				9m; 10y (subnational)	≥60y; risk groups		risk groups; indigenous populations	special groups	
CHI		1y	6y			7y							≥60y; risk groups; pregnant women; health workers			
COL		1y	5y			WCBAs					1y; population in at-risk areas;	6m-23m; ≥60y				
COR		15m	6-7y			7y	every 10y					risk groups; 6m -<8y with risk factors; ≥65y		15m		
CUB		1y	6y			15y							≥65y; children <24y w/chronic diseases			
DOR		1y	5y			7y	10y	every 10 years					6m-3y; >65y; persons w/chronic diseases			
ECU		12-23m	6y			11y	WCBAs				>1y at risk areas	6-23m ; ≥65y		6y; 10y		
ELS		1y	4y			≥10y	every 10y; pregnant women					6m-23m; >60y; risk groups				
GUT		1y				WCBAs						>60y				
HAI					9m	WCBAs										
HON		1y				11-12y	pregnant women; WCBAs					6-35m ; ≥60y; risk groups				
MEX*		1y	6y			11-12y (Grade 6)	WCBAs					6m-35m; ≥60y and risk groups		1y (social security)		
NIC		1y				10y	20y	WCBAs					6-4y; >65y with chronic disease; health workers			
PAN		1y	4y			6y+	WCBAs				>1y population in at-risk areas	every 10y	6m-5y; >60y; risk groups		12m	18m
PAR		1y	4y			10y	WCBAs				1y; ≥1y in population in border areas		6-35m; >60; other risk groups			
PER		1y	4y			WCBAs					15m		6-23m ; ≥65y			
URU		1y	5y			12y	every 10y						6-29m; 65y+; risk groups;	1y	15m	21m
VEN		1y	5y			10y	WCBAs				1y*		6m; ≥60y; risk groups			

NB: Newborn

WCBA(s): Women of Childbearing Age

† Yellow Fever vaccine administered to travellers to endemic countries not included

* PAN: Introduced Hepatitis A vaccines in May 2006.

* VEN: Yellow fever vaccine administered to recruits and travelers to endemic areas.

DISCLAIMER: Table not yet approved by countries

Updated: 28 September 2011



Source: Country reports through PAHO-WHO/UNICEF Joint Reporting Form (JRF)



2010 Immunization Schedule for Selected Vaccines - Latin American Countries

Country	Rotavirus			Vitamin A						Pneumococcal_Conjugate					Meningitis BC		Meningitis AC		Meningitis C					
	1	2		Number of Doses						1	2	3	4	5	1	2	1	2						
ARG*												risk groups												
BOL*	2m	4m		6-11m	1-4y																			
BRA*	2m	4m										2m	4m	6m	15m						3m	5m	>1	
CHI												2m*	4m*	6m*										
COL	2m	4m										2m*	4m*	6m*										
COR												2m	4m	6m	15m									
CUB																	3m	5m						
DOR																								
ECU	2m	4m		6m	12m	18m	24m	30m	36m			2m*	4m*	6m*			risk groups							
ELS	2m	4m										2m	4m	6m	12m									
GUT	2m	4m	6m*	6m	12m	18m	24m	30m	36m	42m	48m	54m												
HAI				6m	10m	14m	18m	30m	42m															
HON	2m	4m		6m	12m	18m	24m	30m	36m			intro in 2011												
MEX*	2m	4m		6-59m									2m	4m	12m									
NIC	2m	4m	6m	6m	1y	2y	3y	4y				(intro in 2011)												
PAN	2m	4m		6m	12m	18m						2m*	4m*	6m*	12-15m*									
PAR	2m	4m																						
PER	2m	4m		6m*	12m*	23m*						3m	5m	12m										
URU												2m	4m	12m										
VEN	2m	4m																						

NB: Newborn
WCBA(s): Women of Childbearing Age

- * ARG: Argentinian hemorrhagic fever for groups at risk
- * BRA: Year of introduction of pneumococcal conjugate is 2010
- * COL: Pneumococcal conjugate given to population at risk
- * CHI: Pneumococcal conjugate given to risk groups
- * GUT: originally a 3-dose vaccine used
- * MEX: SNS-Semana Nacional de Salud; Pneumo_conj given
- * PER: At subnational

DISCLAIMER: Table not yet approved by countries
Updated: 28 September 2011



Source: Country reports through PAHO-WHO/UNICEF Joint Reporting Form (JRF)



2010 Immunization Schedule for Selected Vaccines - Latin American Countries

Country	Typhoid Fever			Tdap	Pneumo_Polysaccharide (v23)	HPV
	1	2	3	1	1	1
ARG*	armed forces starting 2009			11y; health workers	risk groups	
BOL*						
BRA*	risk groups				risk groups	
CHI					65y+	
COL						
COR				risk groups		
CUB	10y	13y	16y			
DOR					>65y	
ECU					older adults >65y and chronic disease	
ELS					with chronic disease	
GUT						
HAI						
HON						
MEX*					≥65y	12y; risk municipalities
NIC					≥50y	
PAN				11y	>60y	girls 10y (3 doses)
PAR					risk groups	
PER						
URU					≥65y	
VEN					≥60y	

DISCLAIMER: Table not yet approved by countries
Updated: 28 September 2011



Source: Country reports through PAHO-WHO/UNICEF Joint Reporting Form (JRF)

