Paraguay (Ages 13-15)	youth tobacco
Global Youth Tobacco Survey (GYTS)	tobacco
	survey

The Paraguay GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Paraguay could include in a comprehensive tobacco control program.

The Paraguay GYTS was a school-based survey of students in  $7^{th}$ ,  $8^{th}$  and  $9^{th}$  grade conducted in 2008.

#### Prevalence

21.6% of students had ever smoked cigarettes (Boy = 26.0%, Girl = 17.6%)

16.7% currently use any tobacco product (Boy = 20.8%, Girl = 12.9%)

8.3% currently smoke cigarettes (Boy = 11.3%, Girl = 5.5%)

10.3% currently use other tobacco products (Boy = 12.4%, Girl = 8.4%)

14.2% of never smokers are likely to initiate smoking next year

#### **Knowledge and Attitudes**

10.7% think boys and 7.9% think girls who smoke have more friends 7.8% think boys and 5.2% think girls who smoke look more attractive

### Access and Availability - Current Smokers

17.2% usually smoke at home

32.8% buy cigarettes in a store

79.5% who bought cigarettes in a store were NOT refused purchase because of their age

#### Exposure to Secondhand Smoke (SHS)

32.5% live in homes where others smoke in their presence

55.3% are around others who smoke in places outside their home

85.7% think smoking should be banned from public places

64.4% think smoke from others is harmful to them

31.6% have one or more parents who smoke

6.8% have most or all friends who smoke

#### **Cessation - Current Smokers**

59.0% want to stop smoking69.0% tried to stop smoking during the past year73.6% have ever received help to stop smoking

#### Media and Advertising

85.0% saw anti-smoking media messages, in the past 30 days
89.0% saw pro-cigarette ads on billboards, in the past 30 days
88.0% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
11.8% have an object with a cigarette brand logo

15.1% were offered free cigarettes by a tobacco company representative

## School

79.4% had been taught in class, during the past year, about the dangers of smoking 48.6% had discussed in class, during the past year, reasons why people their age smoke 69.7% had been taught in class, during the past year, the effects of tobacco use

# Highlights

A two-stage cluster sample design was used to produce

representative data for Paraguay. At the first stage, schools

At the second stage, classes were randomly selected and all

students in selected classes were eligible to participate. The

school response rate was 98.0%, the class response rate was

participated in the Paraguay GYTS.

100.0%, the student response rate was 88.4%, and the overall

response rate was 86.7%. A total of 4,524 students aged 13-15

were selected with probability proportional to enrollment size.

- More than 1 in 10 students currently use any form of tobacco; 8.3% of the students currently smoke cigarettes; 10.3% currently use some other form of tobacco.
- SHS exposure is moderate one third of students live in homes where others smoke, and over half are exposed to smoke around others outside of the home; 3 in 10 students have at least one parent who smokes.
- More than 6 in 10 students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Six in 10 current smokers want to stop smoking.
- 15.1% of students were offered a free cigarette by a tobacco company representative.
- Over 8 in 10 of the students saw anti-smoking media messages in the past 30 days; nearly 9 in 10 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.