

Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory with Autochthonous Transmission

2013-2014 (to week noted)

Cumulative cases

Epidemiological Week / EW 25 (Updated 20 June 2014)

Country/Territory	Week ^a	Cases ^b		Incidence Rate ^c	Deaths	Population ^d
		Suspected	Confirmed	incidence Rate	Deatils	X 1000
Latin Caribbean						
Dominican Republic	Week 25	89,720	18	862.5	3	10,404
French Guiana [*]	Week 24	0	390	156.6	0	249
Guadaloupe	Week 24	40,400	1,328	8,954.5	3	466
Haiti	Week 23	11,802	14	114.5	0	10,317
Martinique	Week 24	35,000	1,515	9,038.4	12	404
Puerto Rico	Week 22	0	23	0.6	0	3,688
Saint Barthelemy	Week 24	620	142	8,561.8	0	9
Saint Martin (French part)	Week 24	3,430	793	11,832.4	3	36
Subtotal		180,972	4,223	724.2	21	25,573
Non-Latin Caribbean						
Anguilla	Week 13		33	206.3	0	16
Antigua and Barbuda	Week 20		4	4.4	0	90
Dominica	Week 20	2,363	129	3,413.7	0	73
Guyana	Week 23		16	2.0	0	800
Saint Kitts and Nevis	Week 21		22	43.1	0	51
Saint Lucia	Week 23	214	30	149.7	0	163
Saint Vincent and the Grenadines	Week 23	212	67	270.9	0	103
Sint Maarten (Dutch part) [≡]	Week 12		123	307.5	0	40
Turks and Caicos Islands [%]	Week 25	0	6	12.5	0	48
Virgin Islands (UK)	Week 19	0	20	62.5	0	32
Virgin Islands (US)	Week 25		3	2.9	0	105
Subtotal		2,789	453	213.1	0	1,521
TOTAL		183,761	4,676	695.5	21	27,094

NOTES

Data source: Cases reported by IHR NFPs to PAHO/WHO and/or through Member States websites or official news publication

Includes two imported cases

^a Epidemiological Week for which information is available.

b Suspected case: patient with acute onset of fever >38°C (101°F) and severe arthralgia or arthritis not explained by other medical conditions, and who resides or has visited epidemic or endemic areas within two weeks prior the onset of the symptoms. Confirmed case: a suspected case with any specific CHIK test (viral isolation, RT-PCR, Ig M, or four-fold increase of CHIK specific antibodies titers) - PAHO/CDC confirmed case definition available at www.paho.org/chikungunya.

c Incidence / 100,000 pop.

^d PAHO/WHO. Basic Indicators, 2013. http://www.paho.org/hq/index.php?option=com_docman&task=doc_download&gid=23535&Itemid=270&lang=en, except St. Barthelemy ("Populations légales 2009 entrent en vigueur le 1er janvier 2012." Recensement de la population. INSEE. Retrieved 2 January 2014).

^{*} Includes 86 (22%) imported cases.

E Data published by Le point épidémiologique, N°11, 2014.

[%] Includes four imported cases



Number of Imported Cases of Chikungunya Fever in the Americas, by Country or Territory

Weekly and Cumulative cases

Epidemiological Week / EW 25 (Updated 20 June 2014)

	ounc 201+)					
Country/Territory	Week ^a	Weekly Imported Cases	Cumulative Imported Cases			
North America						
United States of America	Week 25	18	57			
Central America Isthmus						
Panama	Week 21	2	2			
Latin Caribbean						
Cuba	Week 25	1	6			
Andean Area						
Venezuela	Week 25	4	6			
Brazil	Week 25	4	11			
Southern Cone						
Chile	Week 22	1	2			
Non-Latin Caribbean						
Aruba	Week 06	1	1			
Barbados	Week 23	1	1			
Suriname	Week 25	1	1			
TOTAL			87			

NOTES

Data source: Cases reported by IHR NFPs to PAHO/WHO and/or through Member States websites or official news publication

^a Epidemiological Week for which information is available.

^{*} Only imported cases in countries or territories **without** authoctonous trasnmsision; full epidemiology report on "Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory with Autochthonous Transmission"