Antigua and Barbuda 2009 (Ages 13-15)	global
Global Youth Tobacco Survey (GYTS)	youth tobacco
FACT SHEET · · · · · · · · · · · ·	survey

The Antigua and Barbuda GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Antigua and Barbuda could include in a comprehensive tobacco control program.

The Antigua and Barbuda GYTS was a school-based survey of students in Grade 6, Grade 7/Form 1, Grade 8/Form 2, Grade 9/Form 3, and Form 4 conducted in 2009.

A two-stage cluster sample design was used to produce representative data for Antigua and Barbuda. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. A total of 1,795 students participated in the Antigua and Barbuda GYTS of which 1,068 were ages 13 to 15 years. The overall response rate of all students surveyed was 85.5%.

Prevalence

23.2% of students had ever smoked cigarettes (Boys = 23.2%, Girls = 21.4%)

- 20.1% currently use any tobacco product (Boys = 24.3%, Girls = 15.9%)
- 5.2% currently smoke cigarettes (Boys = 5.2%, Girls = 4.3%)
- 17.6% currently use other tobacco products (Boys = 22.1%, Girls = 14.0%)
- 12.8% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

33.1% think boys and 18.6% think girls who smoke have more friends 7.7% think boys and 5.4% think girls who smoke look more attractive

Access and Availability - Current Smokers

44.9% usually smoke at home 31.6% buy cigarettes in a store

Exposure to Secondhand Smoke (SHS)

- 26.7% live in homes where others smoke in their presence
- 47.5% are around others who smoke in places outside their home
- 77.1% think smoking should be banned from public places
- 68.2% think smoke from others is harmful to them

23.3% have one or more parents who smoke

4.6% have most or all friends who smoke

Cessation - Current Smokers

65.7% have ever received help to stop smoking

Media and Advertising

- 66.9% saw anti-smoking media messages, in the past 30 days
- 35.1% saw pro-cigarette ads on billboards, in the past 30 days
- 31.5% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 11.7% have an object with a cigarette brand logo
- 8.7% were offered free cigarettes by a tobacco company representative

School

37.9% had been taught in class, during the past year, about the dangers of smoking 24.3% had discussed in class, during the past year, reasons why people their age smoke 36.6% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 20.1% of students currently use any form of tobacco; 5.2% currently smoke cigarettes; 17.6% currently use some other form of tobacco
- SHS exposure over onequarter of the students live in homes where others smoke, and 47.5% of students are exposed to smoke around others outside of the home; nearly one-quarter of the students have at least one parent who smokes
- More than two-thirds of students think smoke from others is harmful to them
- Close to eight in 10 students think smoking in public places should be banned
- 11.7% of students have an object with a cigarette brand logo on it
- Two-thirds of the students saw anti-smoking media messages in the past 30 days; over onethird of the students saw procigarette ads on billboards; and three in 10 saw pro-cigarette ads in newspapers or magazines in the past 30 days