Global School-based Student Health Survey

Guatemala (Rest of Country) 2009 Fact Sheet



The **2009 Guatemala (Rest of Country) GSHS** measured alcohol use; dietary behaviors; hygiene; mental health; physical activity; and sexual behaviors.

The 2009 Guatemala (Rest of Country) GSHS was a school-based survey of students in grades primero, segundo, and tercero basico. A two-stage cluster sample design was used to produce data representative of all students in grades primero, segundo, and tercero basico in Guatemala (Rest of Country). At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96%, the student response rate was 86%, and the overall response rate was 83%. A total of 1970 students participated in the Guatemala (Rest of Country) GSHS. Students self-reported their responses to each question on a computer scannable answer sheet. For comparison purposes, only students aged 13- 15 years are included in the analyses for this fact sheet.

Results for students aged 13-15 years	Total	Boys	Girls		
Alcohol Use					
Among students who ever had a drink of alcohol (other than a few sips), the percentage who had their first drink of alcohol before age 14 years	70.6	70.9	70.1		
	(62.2-77.8)	(58.5-80.8)	(60.2-78.4)		
Percentage of students who drank at least one drink containing alcohol on one or more of the past 30 days	10.4	11.0	9.8		
	(8.7-12.4)	(8.3-14.4)	(7.3-13.2)		
Percentage of students who drank so much alcohol that they were really drunk one or more times during their life	7.2	8.2	6.2		
	(5.9-8.8)	(6.0-11.1)	(4.8-7.9)		
Dietary Behaviours					
Percentage of students who were underweight (< -2SD from median for BMI for age and sex)	1.4	1.4	1.4		
	(0.7-2.6)	(0.4-4.4)	(0.9-2.1)		
Percentage of students who were overweight ($> +1$ SD from median for BMI for age and sex)	22.8	21.3	24.7		
	(18.0-28.6)	(16.7-26.7)	(18.6-32.1)		
Percentage of students who were obese (> +2SD from median for BMI for age and sex)	5.2	6.3	3.9		
	(3.1-8.6)	(3.8-10.3)	(2.1-7.1)		
Percentage of students who usually drank carbonated soft drinks one or more times per day during the past 30 days	57.6	56.7	58.1		
	(49.6-65.2)	(48.8-64.4)	(49.4-66.3)		
Hygiene					
Percentage of students who usually cleaned or brushed their teeth less than one time per day during the past 30 days	3.6	4.3	2.7		
	(2.6-5.1)	(3.1-6.1)	(1.7-4.1)		
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the past 30 days	3.2	4.6	1.6		
	(2.3-4.4)	(2.7-7.5)	(0.7-3.4)		
Mental Health					
Percentage of students who ever seriously considered attempting suicide during the past 12 months	12.4	9.1	16.1		
	(9.7-15.6)	(7.8-10.6)	(11.2-22.5)		
Percentage of students who actually attempted suicide one or more times during the past 12 months	13.1	10.2	16.4		
	(10.3-16.5)	(6.5-15.8)	(11.9-22.1)		
Percentage of students who had no close friends	6.3	6.3	5.9		
	(5.1-7.8)	(4.1-9.4)	(3.7-9.3)		
Physical Activity					
Percentage of students who were physically active for a total of at least 60 minutes per day on five or more days during the past seven days	20.7	25.3	15.4		
	(16.7-25.4)	(19.7-31.9)	(12.0-19.6)		
Percentage of students who went to physical education (PE) class on three or more days each week during the school year	26.7	28.7	24.3		
	(23.9-29.7)	(25.2-32.4)	(20.6-28.3)		
Percentage of students who spent three or more hours per day during a typical or usual day doing sitting activities	17.2	16.5	18.4		
	(15.1-19.6)	(13.7-19.8)	(15.3-21.9)		

2009 Page 1

Global School-based Student Health Survey

Guatemala (Rest of Country) 2009 Fact Sheet



Results for students aged 13-15 years	Total	Boys	Girls
Sexual Behaviours			
Percentage of students who ever had sexual intercourse	11.8 (9.5-14.4)	15.8 (12.5-19.7)	7.0 (5.2-9.4)
Among students who ever had sexual intercourse, the percentage who had sexual intercourse for the first time before age 14 years	56.7 (44.2-68.4)	*	*
Among students who ever had sexual intercourse, the percentage who used a condom the last time they had sexual intercourse	57.5 (46.2-68.1)	*	*

^{*} Indicates less than 100 students.

For additional information, please contact:

Lic. Maggie Fischer

Organización Panamericana de la Salud, Guatemala - Rest of Country, fischerm@gut.ops-oms.org

2009 Page 2