Pan American Health Organization World Health Organization

Adolescent and Youth Health - 2017 Country Profile

BARBADOS



EDUCATION PARTICIPATION

(20-24 years old)



ECONOMIC DEVELOPMENT

Source: World Development Indicators (WDIs), The World Bank, 2016.




MORTALITY RATES - Four Leading Causes of Death, by Age Group and Sex (per 100,000 Population)


Source: PAHO Regional Mortality Database, Pan American Health Organization (PAHO/WHO), 2016 Edition. Country latest mortality data available as of 2013. Note: * If there is no cause of death reported for a specific age group and sex, the corresponding bar with zero value is not being represented.

(10-14 years old)

(15-19 years old)



MORTALITY TRENDS - Four Leading Causes of Death, by Sex (per 100,000 Population)





Adolescent and Youth Health - 2017 Country Profile BARBADOS



SEXUAL AND REPRODUCTIVE HEALTH MALE FEMALE ΤΟΤΑΙ **Trends in Adolescent Fertility Rate** Sexual Behavior in Adolescents (Births per 1,000 women ages 15-19 years old), 1980-2015 (age group: 13-15 years old) Source: United Nations Population Divisio Source: 2011 Global School Health Survey (GSHS), WHO World Population Prospects: The 2015 Revision 23.1% 44.0% 100 Students who ever had sexual intercourse 67 33.5% 56.1% 74.3% 50 Among students who ever had 46 47 \sim sexual intercourse, those that the first time was before age 14 yrs 28 68.1% 65.3% Among students who ever had 0 64.6% sexual intercourse, those who 1980-1985 1985-1990 1990-1995 1995-2000 2000-2005 2005-2010 2010-2015 used a condom the last time 64.9% Barbados North America 60% 100% 20% 80% 0% 40% Latin America and the Caribbean World **RISK FACTORS** MALE FEMALE TOTAL GSHS - Global School Health Survey (WHO) GYTS - Global Youth Tobacco Survey (PAHO) OAS - Report on Drug Use in the Americas Sources **Alcohol Use Tobacco Use** 88.7% 17.4% Among students who ever drunk 2011 GSHS 88.6% 11.4% 2013 GYTS Prevalence of current use of any alcohol, those that the first drink was 14.5% 88.7% tobacco product before age 14 yrs 48.0% 58.5% Students who drank at least one drink 2011 GSHS 45.8% Students who reported people 2011 GSHS 55.6% containing alcohol 1+ of the past 30 46.9% smoked in their presence, past 7 days 57.1% days 51.2% 88.2% Among students who ever smoked 54.4% Binge drinking among secondary school 2015 OAS 2011 GSHS 82.2% cigarettes, those that first tried 53.3% students 85,9% cigarettes before age 14 yrs **Overweight/Obesity/Diet Physical Activity** 35.4% Students who went to physical 74.9% 31.2% 2011 GSHS Students who usually drank carbonated 71.5% education (PE) class 3+ days weekly in 2011 GSHS 33.8% soft drinks 1+ times per day the past 30 the school year days 60.4% 32.0% Students who spent 3+ hours per day 2011 GSHS 69.6% 31.8% Students who were overweight (> +1SD 2011 GSHS doing sitting activities 64.9% from median for BMI for age and sex) 31.9% 34.5% 13.9% Students physically active at least 60 23.3% 2011 GSHS Students who were obese (> +2SD from 2011 GSHS 14.6% minutes daily on 5+ days the past 7 29.1% median for BMI for age and sex) 14.2% days **Drug Use Violence and Unintentional Injuries** Among students who ever used drugs. 46.4% 2011 GSHS those that first used drugs before age 2011 GSHS 35.9% Students seriously injured 1+ times 14 vrs 41.3% during the past 12 months Prevalence of cocaine use among 2015 OAS 47.9% secondary school students (lifetime) Students who were in a physical fight 2011 GSHS 28.3% 26.0% 1+ times the past 12 months 38.4% Prevalence of marijuana use among 2015 OAS 18.6% secondary school students (lifetime) 15.4% 11.0% 18.6 Students who were bullied 1+ days 2011 GSHS Prevalence of inhalants use among 22.5% 2015 OAS during the past 30 days 13.3% secondary school students (lifetime)

C Pan American Health Organization (PAHO/WHO). All rights reserved.

www.paho.org