Mexico - Oaxaca (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Mexico - Oaxaca GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Mexico - Oaxaca could include in a comprehensive tobacco control program.

The Mexico - Oaxaca GYTS was a school-based survey of students in grades 1, 2 and 3 conducted in 2003.

A two-stage cluster sample design was used to produce representative data for Mexico - Oaxaca. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96.0%, the class response rate was 100.0%, the student response rate was 93.5%, and the overall response rate was 89.7%. A total of 1,380 students aged 13-15 participated in the Mexico - Oaxaca GYTS.

Prevalence

45.2% of students had ever smoked cigarettes (Boy = 50.7%, Girl = 39.4%)

19.1% currently use any tobacco product (Boy = 22.4%, Girl = 15.6%)

13.8% currently smoke cigarettes (Boy = 14.5%, Girl = 12.7%)

7.6% currently use other tobacco products (Boy = 10.8%, Girl = 4.4%)

24.5% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

21.6% think boys and 32.7% think girls who smoke have more friends 11.3% think boys and 27.1% think girls who smoke look more attractive

Access and Availability - Current Smokers

18.1% usually smoke at home

43.2% buy cigarettes in a store

68.8% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

23.0% live in homes where others smoke in their presence

41.9% are around others who smoke in places outside their home

90.3% think smoking should be banned from public places

79.2% think smoke from others is harmful to them

27.1% have one or more parents who smoke

53.3% have most or all friends who smoke

Cessation - Current Smokers

57.0% want to stop smoking

59.0% tried to stop smoking during the past year

70.8% have ever received help to stop smoking

Media and Advertising

83.1% saw anti-smoking media messages, in the past 30 days

84.7% saw pro-cigarette ads on billboards, in the past 30 days

73.9% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

10.3% have an object with a cigarette brand logo

7.4% were offered free cigarettes by a tobacco company representative

School

59.5% had been taught in class, during the past year, about the dangers of smoking 33.7% had discussed in class, during the past year, reasons why people their age smoke 48.9% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 19.1% of students currently use any form of tobacco; 13.8% currently smoke cigarettes; 7.6% currently use some other form of tobacco.
- SHS exposure is high close to one-quarter of the students live in homes where others smoke, and 2 in 5 students are exposed to smoke around others outside of the home; nearly 3 in 10 students have at least one parent who smokes.
- Eight in 10 students think smoke from others is harmful to them.
- Nine in 10 students think smoking in public places should be banned.
- Close to 3 in 5 current smokers want to stop smoking.
- One in 10 students has an object with a cigarette brand logo on it.
- Over 4 in 5 students saw antismoking media messages in the past 30 days; more than threequarters of the students saw pro-cigarette ads on billboards in newspapers or magazines in the past 30 days.