



## Emerging and Reemerging Infectious Diseases, Region of the Americas

Vol. 1, No. 10—12 September 2003

### Yellow Fever Situation in the Americas

As of 10 September 2003, a total of 184 cases of confirmed jungle yellow fever have been reported to the Pan American Health Organization, with 85 deaths. This represents the highest number of cases reported in the Region since 1999.

**Table 1: Jungle Yellow Fever, Reported Cases and Deaths  
(by country, Region of the Americas, as of 10 September 2003)**

Country	Cases	Deaths
Bolivia	6	4
Brazil	62	23
Colombia	88	40
Peru	15	9
Venezuela	13	9
<b>Total</b>	<b>184</b>	<b>85</b>

*Source:* Reports to PAHO by the Ministries of Health of the respective countries.

The outbreak on the Colombian-Venezuelan border, which started in early June 2003, has moved northward. Four cases were identified during Epidemiological Weeks (EW) 33 and 34 (ending on 16 and 23 August, respectively) in the counties of Agustin Codazzi, Becerril and La Jagua de Ibirico, in the Department of Cesar, northeast region of Colombia. These counties are located across the border from the counties of Machiques and Rosario de Perija, in the State of Zulia in Venezuela, which have also reported cases during EW 31 and 32 (ending on 2 and 9 August, respectively).

The urban areas close to the outbreak are infested with *Aedes aegypti*, highlighting the risk of a possible re-urbanization of the disease in the Americas. An active search for febrile-icteric and febrile-icterohemorrhagic cases is being conducted in all affected Departments. The objective is early detection of YF virus circulation, which will make timely implementation of outbreak-control measures possible. In addition, intensive mass-vaccination efforts are underway, aimed at immunizing all residents in the States of Norte Santander and Cesar, in Colombia, as well as the States of Zulia and Tachira in Venezuela. Public-health authorities from both countries have developed joint border activities to cope with the outbreak and to overcome the difficulties of vaccinating areas with displaced populations and guerrilla activity.

To prevent the occurrence of jungle yellow fever outbreaks and the re-urbanization of the disease in the Americas, enzootic countries should implement all the [recommendations](#) of the PAHO Technical Advisory Group on Vaccine Preventable Diseases. (TAG).

*Source:* Ministries of Health of Bolivia, Brazil, Colombia, Peru and Venezuela.