

Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory
2015 (to week noted)
Cumulative cases
Epidemiological Week / EW 30 (Updated as of 31 July 2015)

| Country/Territory | Week ^a | Autochthonous transmission cases ^b | | Imported cases | Incidence Rate ^c | Deaths ^d | Population ^e X 1000 |
|---------------------------------------|-------------------|---|---------------|----------------|-----------------------------|---------------------|-----------------------------------|
| | | Suspected | Confirmed | | | | |
| North America | | | | | | | |
| Bermuda | Week 17 | | | 3 | | | 69 |
| Canada | Week 4 | | | 312 | | | 35,182 |
| Mexico | Week 30 | | 2,646 | 11 | 2.2 | | 118,129 |
| United States of America ^g | Week 30 | | | 249 | | | 320,051 |
| Subtotal | | 0 | 2,646 | 575 | 0.6 | 0 | 473,431 |
| Central American Isthmus | | | | | | | |
| Belize | Week | | | | 0.0 | | 332 |
| Costa Rica | Week 20 | | 142 | | 2.9 | | 4,872 |
| El Salvador | Week 27 | 23,811 | 13 | | 375.8 | 0 | 6,340 |
| Guatemala | Week 21 | 7,342 | 522 | | 50.8 | 1 | 15,468 |
| Honduras | Week 26 | 49,162 | 5 | | 607.1 | 0 | 8,098 |
| Nicaragua | Week 12 | 17,946 | 2,235 | | 331.9 | | 6,080 |
| Panama | Week 27 | | 13 | 13 | | | 3,864 |
| Subtotal | | 98,261 | 2,930 | 13 | 224.6 | 1 | 45,054 |
| Latin Caribbean | | | | | | | |
| Cuba | Week | | | | 0.0 | | 11,266 |
| Dominican Republic | Week 22 | 63 | | | 0.6 | | 10,404 |
| French Guiana | Week 30 | 6,380 | 1,756 | | 3,267.5 | 2 | 249 |
| Guadeloupe | Week 9 | 150 | | | 32.2 | | 466 |
| Haiti | Week | | | | | | 10,317 |
| Martinique | Week 9 | 320 | | | 79.2 | | 404 |
| Puerto Rico ^f | Week 27 | 575 | 103 | | 18.4 | | 3,688 |
| Saint Barthelemy | Week 18 | 317 | | | 3,561.8 | | 9 |
| Saint Martin (French part) | Week 18 | 600 | | | 1,681.1 | | 36 |
| Subtotal | | 8,405 | 1,859 | 0 | 27.9 | 2 | 36,839 |
| Andean Area | | | | | | | |
| Bolivia | Week 13 | 143 | 916 | 1 | 9.9 | | 10,671 |
| Colombia [®] | Week 28 | 307,658 | 864 | | 638.5 | 37 | 48,321 |
| Ecuador | Week 30 | 27,156 | 3,297 | 95 | 193.5 | 2 | 15,738 |
| Peru | Week 24 | 11 | 7 | 27 | 0.1 | | 30,376 |
| Venezuela | Week 17 | 12,482 | 296 | | 42.0 | | 30,405 |
| Subtotal | | 347,450 | 5,380 | 123 | 260.4 | 39 | 135,511 |
| Southern Cone | | | | | | | |
| Argentina | Week 28 | | | 30 | | | 41,446 |
| Brazil | Week 26 | 8,251 | 120 | 7 | 4.2 | | 200,362 |
| Chile | Week 28 | | | 6 | | | 17,620 |
| Paraguay | Week 27 | 1,854 | 840 | | 39.6 | | 6,802 |
| Uruguay | Week | | | | | | 3,419 |
| Subtotal | | 10,105 | 960 | 43 | 4.1 | 0 | 269,649 |
| Non-Latin Caribbean | | | | | | | |
| Anguilla | Week 11 | 28 | 3 | | 193.8 | | 16 |
| Antigua and Barbuda | Week 8 | 16 | 0 | | 17.8 | | 90 |
| Aruba | Week 14 | 66 | 686 | | 689.9 | | 109 |
| Bahamas | Week 17 | | 10 | 1 | 2.7 | | 377 |
| Barbados | Week 19 | 290 | 15 | | 105.5 | | 289 |
| Cayman Islands | Week 8 | 67 | 1 | | 125.9 | | 54 |
| Curacao | Week | | | | | | 147 |
| Dominica | Week 5 | 8 | | | 11.0 | | 73 |
| Grenada | Week | | | | | | 110 |
| Guyana | Week 2 | 5,310 | 29 | | 667.4 | | 800 |
| Jamaica | Week 6 | 299 | | | 10.7 | | 2,784 |
| Montserrat | Week 20 | 13 | 0 | | 260.0 | | 5 |
| Saint Kitts and Nevis | Week | | | | | | 51 |
| Saint Lucia | Week | | | | | | 163 |
| Saint Vincent and the Grenadines | Week 8 | 3 | 2 | | 4.9 | | 103 |
| Sint Maarten (Dutch part) | Week | | | | | | 40 |
| Suriname | Week | | | | | | 539 |
| Trinidad and Tobago | Week 20 | | 83 | | 6.2 | | 1,341 |
| Turks and Caicos Islands | Week | | | | | | 48 |
| Virgin Islands (UK) | Week | | | | | | 32 |
| Virgin Islands (US) | Week 12 | 225 | 102 | | 311.4 | | 105 |
| Subtotal | | 6,325 | 931 | 1 | 99.7 | 0 | 7,276 |
| TOTAL | | 470,546 | 14,706 | 755 | 50.1 | 42 | 967,760 |

NOTES

^a Epidemiological Week for which information is available.

^b **Suspected case:** patient with acute onset of fever >38°C (101°F) and severe arthralgia or arthritis not explained by other medical conditions, and who resides or has visited epidemic or endemic areas within two weeks prior to the onset of the symptoms.
Confirmed case: a suspected case with any specific CHIK test (viral isolation, RT-PCR, Ig M, or four-fold increase of chikungunya specific antibodies titers) - PAHO/CDC confirmed case definition available at www.paho.org/chikungunya.

^c Incidence rate (autochthonous suspected + autochthonous confirmed) / 100,000 pop.

^d Deaths directly or indirectly related to Chikungunya.

^e PAHO/WHO. Basic Indicators, 2013. http://www.paho.org/hq/index.php?option=com_docman&task=doc_download&gid=23535&Itemid=270&lang=en, except St. Barthelemy ("Populations légales 2009 entrent en vigueur le 1er janvier 2012." Recensement de la population. INSEE. Retrieved 2 January 2014).

^f Department of Health data: <http://www.salud.gov.pr/Estadisticas-Registros-y-Publicaciones/Pages/Chikungunya.aspx>

^g The total cases excludes the data from Puerto Rico and US Virgin Islands.

[®] Confirmed autochthonous cases are cumulative from 2015 EW12, as published in <http://www.ins.gov.co/boletin-epidemiologico/Paginas/default.aspx>

Data source: Cases reported by IHR NFPs to PAHO/WHO and/or through Member States websites or official news publication