

## Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory 2017 (to week noted) Cumulative cases

Epidemiological Week / EW 12 (Updated as of 24 March 2017)

		Autochthonous	s transmission cases <sup>b</sup>				Donulation <sup>®</sup>
Country/Territory	Epidemiological Week <sup>a</sup>	Suspected	Confirmed	Imported cases	Incidence Rate <sup>c</sup>	Deaths <sup>d</sup>	Population <sup>e</sup> X 1000
North America		Cuoposiou					X IOO
Bermuda	Week				0.0		71
Canada	Week						36,626
Mexico	Week 10		8	0	0.01	0	130,223
United States of America >	Week 11			3	0.0	0	
Subtotal		0	8	3	0.00	0	493,394
Central American Isthmus							
Belize	Week				0.0		374
Costa Rica	Week 9	99			2.02		4,905
El Salvador	Week 10	123	0	0	1.99	0	-,
Guatemala	Week				0.00		17,005
Honduras	Week		•		0.00		8,304
Nicaragua	Week 8 Week 8	317	8	1	0.13 7.83		6,217 4,051
Panama Subtotal	vveek o	539	8	1	1.16	0	47,023
Latin Caribbean		555	<u> </u>	,	1.10	V	41,023
Cuba	Week	Ī					11,390
Dominican Republic	Week				0.00		10,766
French Guiana	Week				0.00		282
Guadaloupe	Week				0.00		472
Haiti	Week				0.00		10,983
Martinique	Week						396
Puerto Rico	Week 9		5		0.14		3,679
Saint Barthelemy	Week 5	0	0		0.00		7
Saint Martin (French part)	Week 5	0			0.00		32
Subtotal		0	5	0	0.01	0	38,007
Andean Area							
Bolivia	Week 8	242	0		2.19		11,052
Colombia	Week 10	323	5	0	0.67	0	
Ecuador	Week 10		29		0.17		16,625
Peru**	Week 10	289	28		0.99		32,166
Venezuela	Week 8	39	4	0	0.13	0	31,925
Subtotal		893	66	0	0.68	0	140,835
Southern Cone							
Argentina	Week 10	0	0		0.00		44,272
Brazil	Week 7	8,116	2,178		4.87	1	211,243
Chile	Week 3	,	,	1	0.00		18,313
Paraguay*	Week 8	632	0		9.28		6,811
	Week 2	032		1	5.20		3,456
Uruguay	Week Z	0.740	0.470	2	2.05	-	
Subtotal		8,748	2,178		3.85	1	284,095
Non-Latin Caribbean							
Anguilla	Week				0.00		17
Antigua and Barbuda	Week				0.00		95
Aruba Bahamas	Week Week				0.00		115 397
Barbados	Week				0.00		292
Cayman Islands	Week				0.00		58
Curação	Week				3.00		149
Dominica	Week				0.00		74
Grenada	Week				0.00		111
Guyana	Week				0.00		774
Jamaica	Week 9		0		0.00		2,813
Montserrat	Week				0.00		5
Saint Kitts and Nevis	Week				0.00		52
Saint Lucia	Week				0.00		164
Saint Vincent and the Grenadines	Week				0.00		102
Sint Maarten (Dutch part)	Week				0.00		42
Suriname	Week				0.00		552
Trinidad and Tobago	Week				0.00		1,369
Turks and Caicos Islands	Week				0.00		52
Virgin Islands (UK)	Week				0.00		35
Virgin Islands (US) Subtotal	Week	0	0	0	0.0		102 7,370
TOTAL			2,265	6	1.23	0	7,370 1,010,724
	ar 2017 are precented	10,180	2,200	b	1.23		1,010,724
NOTES: only accumulated cases for the year 2017 are presented.							

<sup>a</sup> Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat, World Population Prospects: The 2015 Revision, http://esa.un.org/unpd/wpp/index.htm, July 2015. International Programs Center, Population Division, U.S. Census Bureau. IDB Release Date: December 2013

In EW 5 of 2017, the number of suspected (cumulative) cases changed from 2 to 423, because both suspected and probable cases were included. http://vigisalud.gov.py/boletines/03\_03\_2017\_12\_07\_12\_Boletin-Epidemiologico\_SE-5.pdf

\*\*On March 3, 2017, the number of probable cases were changed from 175 to 170, based on the modification by the Ministry of Health of Peru. On March 17, 2017, the number of confirmed cases have decreased from 28 to 27, based on the modification by the Ministry of Health of Peru: http://www.dge.gob.pe/portal/docs/vigilancia/sala/2017/salaSE09.pdf

f Department of Health data: http://www.salud.gov.pr/Estadisticas-Registros-v-Publicaciones/Pages/Informe-Arboviral.aspx

Data source: Cases reported by IHR NFPs to PAHO/WHO and/or through Member States websites or official news publication. Cases reported by CARPHA for non-Latin Caribbean countries, unless other source specified

Epidemiological Week for which information is available. Changes in the data from week to week should be interpreted by taking into account the differences in surveillance systems.

Suspected case: patient with acute onset of fever >38°C (101°F) and severe arthralgia or arthritis not explained by other medical conditions, and who resides or has visited epidemic or endemic areas within two weeks prior the onset of the symptoms.

Confirmed case: a suspected case with any specific CHIK test (viral isolation, RT-PCR, Ig M, or four-fold increase of chikungunya specific antibodies titers) - PAHO/CDC confirmed case definition available at www.paho.org/chikungunya.

Incidence rate ( autochthonous suspected + autochthonous confirmed) / 100,000 pop.

d Deaths directly or indirectly related to Chikungunya