


Zika-Epidemiological Report Dominican Republic

25 September 2017

Figure 1. Suspected and confirmed Zika virus cases by epidemiological week (EW). Dominican Republic. EW 48 of 2015 to EW 30 of 2017.



Source: Data provided by the Dominican Republic Ministry of Public Health to PAHO/WHO¹

FIRST AUTOCHTHONOUS VECTOR-BORNE CASES

In epidemiological week (EW) 3 of 2016, the detection of the first autochthonous Zika cases of vector-borne transmission was reported by the Dominican Republic International Health Regulations (IHR) National Focal Point (NFP). The first confirmed cases were from the National District and the municipalities of Santo Domingo Norte, Jimani-Independencia, and Santa Cruz-Barahona.

GEOGRAPHIC DISTRIBUTION

Information regarding geographic distribution of Zika cases is only available up to EW 12 of 2017. As of EW 12 of 2017, autochthonous transmission of Zika virus was confirmed in 31 of the 32 provinces of the Dominican Republic.² Although no confirmed case was detected in the province of Elías Piña, the latter has reported suspected cases. The provinces that reported the highest Zika virus incidence rates are San Jose de Ocoa (210 cases per 100,000 population), Independencia (139 cases per 100,000), Distrito Nacional (100 cases per 100,000), Santo Domingo (78 cases per 100,000), and Valverde (73 cases per 100,000) (**Figure 2**).

Pan American Health Organization • <u>www.paho.org</u> • © PAHO/WHO, 2017

¹ Reported to PAHO/WHO by the Dominican Republic IHR NFP on 21 August 2017.


² Dominican Republic Ministry of Public Health. Weekly epidemiological bulletin. EW 12 of 2017. Available at:

http://digepisalud.gob.do/docs/Boletines%20epidemiol%C3%B3gicos/Boletines%20semanales/2017/Bolet%C3%ADn%20Se manal%2012-2017.pdf

Suggested citation: Pan American Health Organization / World Health Organization. Zika - Epidemiological Report Dominican Republic. September 2017. Washington, D.C.: PAHO/WHO; 2017



Figure 2. Confirmed and suspected Zika cases per 100,000 population by province. Dominican Republic. EW 1 of 2016 to EW 12 of 2017.



Source: Data published by the Dominican Republic Ministry of Public Health and reproduced by PAHO/WHO²

TREND

The peak of the Zika outbreak occurred in EW 15 of 2016, when 354 cases were reported (**Figure 1**).¹ Since then, progressively lower numbers of Zika cases were reported in the Dominican Republic, with an average of one case per week being registered in the last 8 reported weeks (EW 23 to EW 30 of 2017).

CIRCULATION OF OTHER ARBOVIRUSES

As of EW 30 of 2017, a total of 729 probable cases of dengue (6 cases per 100,000 population), were reported) were reported in the Dominican Republic.³ In 2016, a total of 6,645 probable cases of dengue (62 cases per 100,000) were reported up to EW 52.

Suggested citation: Pan American Health Organization / World Health Organization. Zika - Epidemiological Report Dominican Republic. September 2017. Washington, D.C.: PAHO/WHO; 2017

Pan American Health Organization • <u>www.paho.org</u> • © PAHO/WHO, 2017

³ PAHO/WHO. Data, Maps and Statistics. Number of reported cases of Dengue and Severe Dengue (SD) in the Americas. Available at: <u>http://www.paho.org/hq/index.php?option=com_topics&view=rdmore&cid=6290&Itemid=40734</u>



In regard to chikungunya, no information on the number of chikungunya cases reported in 2017 is available.⁴ In 2016, a total of 112 suspected cases (one case per 100,000) were registered up to EW 30. In 2015, 67 suspected cases (one case per 100,000) were detected up to EW 28. In 2014, 524,297 suspected and 84 laboratory-confirmed cases (cumulative incidence rate of 5,040 cases per 100,000) were registered up to EW 45.

ZIKA VIRUS DISEASE IN PREGNANT WOMEN

As of EW 30 of 2017, a total of 966 pregnant women suspected for Zika virus infection have been reported by the Dominican Republic public health authorities.¹ Of the total, 271 pregnant women have been laboratory confirmed for Zika virus infection.

ZIKA COMPLICATIONS

ZIKA-VIRUS-ASSOCIATED GUILLAIN-BARRE SYNDROME (GBS)

Since the confirmation of the first cases of Zika in the Dominican Republic in EW 3 of 2016, an increasing trend of Guillain-Barré syndrome (GBS) cases, similar to the trend of Zika cases was observed (**Figure 4**). The GBS cases, along with the Zika cases steadily declined since EW 23 of 2016. As of EW 30 of 2017, a total of 285 GBS cases, including 32 confirmed cases associated with Zika virus infection were reported by the Dominican Republic public health authorities.¹




Figure 4. Zika and GBS cases by EW. Dominican Republic. EW 48 of 2015 to EW 30 of 2017.

Source: Data provided by the Dominican Republic Ministry of Public Health to PAHO/WHO¹

CONGENITAL SYNDROME ASSOCIATED WITH ZIKA VIRUS INFECTION

As of EW 17 of 2017, 85 cases of microcephaly in newborns with laboratory-confirmed for Zika virus infection have been reported by the Dominican Republic public health authorities to PAHO/WHO.^{1, 5}

Suggested citation: Pan American Health Organization / World Health Organization. Zika - Epidemiological Report Dominican Republic. September 2017. Washington, D.C.: PAHO/WHO; 2017

Pan American Health Organization • <u>www.paho.org</u> • © PAHO/WHO, 2017

⁴ PAHO/WHO. Chikungunya: Statistic Data. Number of reported cases of Chikungunya Fever in the Americas. Available at: <u>http://www.paho.org/hq/index.php?option=com_topics&view=readall&cid=5927&Itemid=40931&lang=en</u>

⁵ The difference between the number of reported cases of confirmed congenital syndrome associated with Zika virus infection from 10 August 2017 (93 cases) to 21 August 2017 (85 cases) is due to a change in the criteria for the case



DEATHS AMONG ZIKA CASES

As of EW 12 of 2017, 17 deaths associated with GBS (mentioned above) were reported by the Dominican Republic public health authorities to PAHO/WHO.^{2, 6}

NATIONAL ZIKA SURVEILLANCE GUIDELINES

Surveillance guidelines for clinicians are being implemented. More information is available at:

http://digepisalud.gob.do/docs/Vigilancia%20Epidemiologica/Alertas%20epidemiologica/Zika/Nacio nal/Procedimiento Vigilancia ZIKV 2016 Ene 31.pdf

In February 2016, "Protocolo De Atención Sindrome De Guillain Barré 2016" was published to standardize the required set of guidelines for diagnosis and management of GBS. More information is available at:

http://digepisalud.gob.do/docs/Vigilancia%20Epidemiologica/Gu%C3%ADa%20de%20Atenci%C3 %B3n/Protocolo%20De%20Atenci%C3%B3n%20Sindrome%20De%20Guillain%20Barr%C3%A9% 202016.pdf

On 14 April 2016, a resolution was passed making it mandatory to report GBS, Microcephaly and any other neurological complications associated with Zika virus disease in the Dominican Republic. More information is available at:

http://digepisalud.gob.do/docs/Vigilancia%20Epidemiologica/Reglamentos%20y%20Normas/2016 %20-%20Resoluci%C3%B3n%20No.%20000013%20%20Notificaci%C3%B3n%20De%20Los%20Casos

<u>%20Resoluci%C3%B3n%20No.%20000013%20%20Notificaci%C3%B3n%20De%20Los%20Casos</u> <u>%20De%20Guillain%20Barr%C3%A9.pdf</u>

LABORATORY CAPACITY

Laboratory confirmation is performed at the *Laboratorio Nacional "Dr. Defilló"* from the Dominican Republic Ministry of Public Health by molecular detection (real-time RT-PCR). In addition, they recently established the PCR multiplex system from the U.S. CDC (Trioplex).

INFORMATION-SHARING

Information on Zika is periodically provided by the Dominican Republic IHR NFP to PAHO/WHO and a periodic epidemiological bulletin is published on the Dominican Republic Ministry of Public Health website. At the time of this report, the latest information on Zika virus provided by the Dominican Republic IHR NFP to PAHO/WHO was from EW 30 of 2017 and the latest epidemiological bulletin with available information on the Zika situation was from EW 17 of 2017. This is because the Dominican Republic Ministry of Health website was unavailable at the time of the preparation of this epidemiological report (accessed on 8 September 2017).

Suggested citation: Pan American Health Organization / World Health Organization. Zika - Epidemiological Report Dominican Republic. September 2017. Washington, D.C.: PAHO/WHO; 2017

definition of microcephaly by the Dominican Republic Ministry of Public Health and Social Assistance, which resulted in the retrospective re-classification of cases.

⁶ Dominican Republic Ministry of Public Health. Weekly epidemiological bulletin. EW 3 of 2017. Available at: <u>http://digepisalud.gob.do/docs/Boletines%20epidemiológicos/Boletines%20semanales/2017/Boletín%20Semanal%2003-2017.pdf</u>