Brazil - Sao Luis (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Brazil - Sao Luis GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Brazil - Sao Luis could include in a comprehensive tobacco control program.

The Brazil - Sao Luis GYTS was a school-based survey of students in 7th and 8th grade of elementary school and 1st grade of high school conducted in 2006.

A two-stage cluster sample design was used to produce representative data for Brazil - Sao Luis. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96.0%, the class response rate was 100.0%, the student response rate was 88.9%, and the overall response rate was 85.4%. A total of 1,389 students aged 13-15 participated in the Brazil - Sao Luis GYTS.

Prevalence

29.3% of students had ever smoked cigarettes (Boy = 28.3%, Girl = 29.2%)

14.7% currently use any tobacco product (Boy = 14.4%, Girl = 14.4%)

10.6% currently smoke cigarettes (Boy = 10.3%, Girl = 10.4%)

4.8% currently use other tobacco products (Boy = 5.2%, Girl = 4.2%)

16.3% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

22.3% think boys and 14.2% think girls who smoke have more friends 6.4% think boys and 4.9% think girls who smoke look more attract

Access and Availability - Current Smokers

26.2% usually smoke at home

23.9% buy cigarettes in a store

Exposure to Secondhand Smoke (SHS)

25.4% live in homes where others smoke in their presence

41.9% are around others who smoke in places outside their home

87.5% think smoking should be banned from public places

78.4% think smoke from others is harmful to them

25.9% have one or more parents who smoke

11.2% have most or all friends who smoke

Cessation - Current Smokers

70.3% want to stop smoking

72.5% tried to stop smoking during the past year

69.4% have ever received help to stop smoking

Media and Advertising

83.8% saw anti-smoking media messages, in the past 30 days

67.3% saw pro-cigarette ads on billboards, in the past 30 days

50.5% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

3.6% have an object with a cigarette brand logo

7.8% were offered free cigarettes by a tobacco company representative

School

54.3% had been taught in class, during the past year, about the dangers of smoking

34.1% had discussed in class, during the past year, reasons why people their age smoke

47.5% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 14.7% of students currently use any form of tobacco; 10.6% currently smoke cigarettes; 4.8% currently use some other form of tobacco.
- SHS exposure is high onequarter of the students live in homes where others smoke, and more than 2 in 5 students are exposed to smoke around others outside of the home; 25.9% of students have at least one parent who smokes.
- Eight in 10 students think smoke from others is harmful to them.
- Nine in 10 students think smoking in public places should be banned.
- Seven in 10 current smokers want to stop smoking.
- 3.6% of students have an object with a cigarette brand logo on it
- Over 8 in 10 students saw antismoking media messages in the past 30 days; two-thirds of the students saw pro-cigarette ads on billboards and half saw protobacco ads in newspapers or magazines in the past 30 days.