





%(508'\$



2019 population (thousands) 71  
Life expectancy (years) 81.6

\$GTOWFC KU C UGNH IQXGTPKPI \$TKV  
#OGTKECU +V EQPUKUVU QH UGXGP  
VJCP UOCNNGT KUNCPFU YKVJ C B  
MO 6JG VGTTKVQT[ KU FKXKFGF KPV  
2CIGV 2GODTQMG 5CPF[U 5OKVJoU  
9CTYKEM 6JG ECRKVC CWGF CKPKNYQDTG  
RQRWNCVGF RCTKUJ  
\$GTOWFC KU IQXGTPGF D[ VJG 9GUVO  
FGOQETCE[  
  
+P VJG RQRWNCVKQP R[TCOKF JO  
UVCVKQPCT[ DGNQY VJG CIGUQMCVKGP  
GZRGVCP[E[ CV DKTUJ KP YCU  
YQOGP  
  
6JG VGTTKVQT[ JCU QPG QH VJG YQTI  
ECRKVC ITQUU FQOGUVKE RTQFWEV  
6JG RTKPEKRCN GEQPQOKE UGEVQTU  
EQPUVTWEVKQP

7+( ' ,6(\$6( %85'(1 \$))(&7,1\* 0(17\$/ +(\$/7+  
0HQVQB XURV XRELVVGL FHRDQCKU FLGHE DXVHR DGOVDELOLW\ DGM  
\HDUV '\$/<V DQG RI DOO \HDUV OLYHG ZLWK GLVDELOLW\



)LJXUH 'LVWULEXWLRQ RI </'V ZLWK D IREFXVHV B HGWZHHQ FKH QHQRULG DO  
VXEVDQFH XVH GLVRUGHUV DQG VHOI KDUP DQG VRPDWLF V\PSWRP GLVRUGHUV

7+( %85'(1 \$))(&7,1\* 0(7+\$\$/526\$/( /,) (7,0(  
)LJ VKRZV WKH FKDQJHV LQ GLVHDVH EXUGHQ DFURVV DJ  
EHWZHHQ DQG \HDUV ROG DQG ZLOO UHPDLQ DERYH  
WKH WRWDO EXUGHQ EHWZHHQ DQG \HDUV RI DJH E\ I  
)LJ IREFXVHV H[FOXVLYHO\ RQ WKH EXUGHQ UHVXOWLQJ I  
DXWLVP DQG HSLOHSV\ %HWZHHQ DQG \HDUV  
DQG KHDGDFKHV rLQFOXGLQJ PLJUDLQH DQG WHQVLRQ W\S  
\$URXQG \HDUV RI DJH D SDWWHUQ HPHUJHV WKDW ZLOO  
DQ[LHW\ GHSUHVLRQ VHOI KDUP DQG VRPDWLF V\PSWRP  
GLVRUGHUV GXH WR DOFRKRO DQG VHYHUH PHQWDC  
HOGHUO\ VXIIHU PRVWO\ IURP QHXURFRJQLWLYH GLVRUGHU  
DURXQG \HDUV ROG DQG UHPDLQV DERYH DIWHU \HE

Figure 3. Burden of disease, by disease group and age





Figure 4. Burden of disease, by MNSS and age



### THE BURDEN AFFECTING MENTAL HEALTH IN MEN AND WOMEN

The top three disorders in terms of disability-adjusted life-years –accounting for 30 to 55% of total MNSS burden- are not the same for men and women: While men are mostly affected by headaches, alcohol and drug use disorders, women are mostly affected by headaches, anxiety and depressive disorders.

Men		Women	
Disorder	DALYs per 100 000	Disorder	DALYs per 100 000
MNSS (all)	3842	MNSS (all)	3872
Headache disorders	525	Headache disorders	940
Alcohol use disorders	418	Anxiety disorders	540
Drug use disorders	358	Depressive disorders	539
Depressive disorders	356	Somatic symptom disorder with prominent pain	384
Alzheimer's disease and other dementias	346	Alzheimer's disease and other dementias	311

### Conclusions:

Considering these estimates, primary care providers should receive training and tools to prioritize detection and treatment or referral for the common disorders highlighted above for each age-group and sex. For the severe disorders –such as autism, schizophrenia, bipolar disorder and Alzheimer's- as well as for severe, co-morbid, or complex presentations of other disorders –e.g. depression during pregnancy, substance use in public service professions, etc.- primary care providers and families need access to adequate supports, such as:

- Referral and/or supervision platforms that allow for continued treatment in the community, including the use of digital technology to increase access to distant geographically concentrated resources.
- Emergency, inpatient, and residential services for the management of high-risk acute situations and high-need patients. These services should be community-based as much as possible, including for crisis management, inpatient treatment in general hospitals, supported housing, and residential services.