

## Situation Report on Mpox Multi-Country Outbreak Response - Region of the Americas

Issue 3/2024, published 25 November 2024

### Contents:

- Highlights
- Region of the Americas – An Epidemiological Overview
- PAHO/WHO Response per Pillar

## MPOX SITUATION IN NUMBERS

### Region of the Americas

As of 31 October 2024

(16:00 EST)

### Total

(13 May 2022 – 31 October 2024)

**66,368**

Confirmed cases

**147**

Deaths

**31**

Countries with confirmed cases

Males – 59,159/61,720 cases  
(95.9%)

Children <18 years –  
7,704/65,483 cases (11.8%)

MSM (Men who have Sex with  
Men) – 14,992/20,767 cases  
(72.2%)

Concurrent HIV Infection –  
13,191/22,496 cases (58.6%)

Healthcare Workers –  
1,265/24,945 cases (5.1%)

Information is updated from Monday to  
Friday by 18:00 GTM-5, at:

[Mpox \(https://shiny.paho-phe.org/mpox/\)](https://shiny.paho-phe.org/mpox/)

## Highlights

Between 2022 and 31 October 2024, a total of 115,101 confirmed cases of mpox have been reported globally. The Region of the Americas (57.6%) contributes the largest proportion of cases, followed by the European (24.7%) and African Regions (12%) [1].

In the Region of the Americas, as of 31 October 2024, 66,368 confirmed cases of Mpox, including 147 deaths, were reported in 31 countries and territories. The highest proportion of cases was recorded in 2022 (57,677 cases, 87%), with a downward trend in 2023 (4,043 cases, 6%) and 2024 (4,648 cases, 7%) (**Figure 1**).

The first case of mpox clade I in the region was identified in the United States of America [2].

## Region of the Americas - An Epidemiological Overview

As of 31 October 2024, the North American subregion reports the highest burden of Mpox cases, with 40,239 cases and 98 deaths (United States with 34,187 cases and 63 deaths, Mexico with 4,184 cases and 35 deaths, and Canada with 1,868 cases) reported up to EW 44 2024. The South American subregion has the next highest proportion of cases (24,941 cases and 44 deaths), followed by Central America (1,022 cases and 3 deaths), and the Caribbean and Atlantic Ocean Islands (166 cases and 2 deaths).

In 2024, a total of 4,221 Mpox cases and 6 deaths have been reported in 14 countries: Argentina (n=87 cases), Bolivia (Plurinational State of) (n=1 case), Brazil (n=1,490 cases), Canada (n=323 cases), Chile (n=10 cases), Colombia (n=131 cases), Costa Rica (n=1 case), Dominican Republic (n=8 cases),


[1] World Health Organization. 2022-24 Mpox (Monkeypox) Outbreak: Global Trends. Geneva: WHO; 2024. [cited 18 November 2024]. Available from: [https://worldhealthorg.shinyapps.io/mpox\\_global/](https://worldhealthorg.shinyapps.io/mpox_global/)

[2] A case of mpox clade I was reported by the United States of America on 16 November 2024. This case has not been included in the case counts in this situation report, which are as of 31 October 2024.

Ecuador (n=24 cases), Guatemala (n=1 case), Mexico (n=105 cases and 1 death), Panama (n=5 cases), Peru (n=90 cases and 2 deaths), the United States (n=2,371 cases and 3 deaths), and Uruguay (n=1 case) (Figure 1).

Most of the cases reported in the Americas Region were identified through human immunodeficiency virus (HIV) care services, sexual health services, or primary and/or secondary health care facilities, involving mainly but not exclusively, men who have sex with men (MSM). The United States of America has reported the identification of the first case of mpox clade I in the region, in a person with recent travel history to East Africa. The individual has been isolated at home, with improving symptoms, and is not receiving specific treatment for mpox. The U.S. CDC is working to identify and follow up with potential contacts [3].

**Figure 1.** Confirmed cases of Mpox by epidemiological week of onset symptoms/notification. Americas Region, as of 31 October 2024.



Figures are subject to change based on the latest data reported by countries/territories

**Source:** Adapted from Pan American Health Organization. Mpox case board – Americas Region. Washington, D.C.: PAHO; 2024 [cited 18 November 2024]. Available from: <https://shiny.paho-phe.org/Mpox/> and from data reported by the IHR National Focal Points to PAHO/WHO.

[3] Pan American Health Organization. Epidemiological Alert Detection of clade I mpox in the Americas Region. 19 November 2024. Washington, D.C.: PAHO; 2024. Available from: <https://www.paho.org/en/documents/epidemiological-alert-detection-clade-i-mpox-americas-region-19-november-2024>

## PAHO/WHO Response per Pillar

### Coordination

PAHO continues to strengthen coordination efforts with Ministries of Health of Member States by supporting epidemiological surveillance, case management, lab diagnosis, community engagement, and risk communication.

### Surveillance

PAHO has been working in close collaboration with local health authorities to help strengthen epidemiological surveillance for Mpox in countries. PAHO, in collaboration with Ministries of Health reviews the situation of Mpox in countries and supports organization of workshops aimed to strengthen the national response in the management and surveillance of Mpox, review infection prevention and control measures. Efforts to provide technical cooperation on surveillance and response to Mpox outbreaks are also being undertaken.

The Organization continued to update the Mpox cases dashboard ([Mpox \(https://shiny.paho-phe.org/mpox/\)](https://shiny.paho-phe.org/mpox/)) and disseminate its use among Member States. It was developed to facilitate data visualization, analysis, and follow-up. The tool is available in English, French, Portuguese, and Spanish. Information is collected through the IHR National Focal Point (NFP) channels and publicly available data from ministries of health.

### Laboratory

PAHO continues efforts to strengthen laboratory capacity in Member States for the rapid detection and diagnosis of Mpox, including procuring equipment, laboratory materials, and reagents.

The organization also provided technical support to the implementation of the Mpox virus detection by PCR, through the provision of supplies, and sharing and reviewing available protocols. Routine meetings are held with staff from laboratories in the Region to review data, test results, troubleshoot, and follow-up on any events in the respective countries.

PAHO has published and updated the Laboratory Guidelines for the Detection and Diagnosis of Monkeypox Virus Infection.

### Clinical Management and Infection Prevention and Control (IPC)

Clade Ib is expected to produce more morbidity and mortality than Clade II. Most of the deaths associated to Mpox were among individuals with advanced HIV infection, unaware of their status or disengaged from care. Therefore, all individuals with lesions suspected to be Mpox should be offered HIV test to be able to start antiretroviral treatment as soon as possible.

PAHO is working with clinicians in Member States to learn and disseminate information on clinical features, diagnostic challenges, and clinical management practices of suspected and confirmed Mpox infections.

The Organization is continuously evaluating IPC interventions that can prevent transmission of Mpox to health care workers in occupational settings in countries in the Region. PAHO routinely participates in meetings with WHO to define the need to update the management guide for cases, and guidelines for infection control and prevention.

Webinars are periodically held to disseminate IPC and clinical management recommendations for persons with Mpox, including [home care](#) of uncomplicated cases.

The [WHO Clinical Platform for Mpox](#) collects anonymized data to understand the clinical features and outcomes of Mpox. Guidance documents for [clinical management and infection prevention and control](#), are being updated.


WHO has launched a call for Expressions of Interest to receive a donation of tecovirimat for use under the Revised MERUI protocol. Countries interested in receiving this drug should contact the local PAHO office for more details. A new [Atlas of mpox lesions](#) has been published to harmonize the assessments of lesions and improve the quality of the collected data.

## Vaccination

During the 2022-2023 period, 13 countries in the region acquired vaccines through the Revolving Fund (RF), as part of their Mpox prevention and control plans.

It is important that countries update their Mpox vaccination plans as part of the national response plan, considering the epidemiological scenario and permanent recommendations, which aim to advance Mpox prevention and control in accordance with the WHO Strategic Framework 2024-2027. This vaccination plan should be based on the most up-to-date recommendations of the WHO Strategic Advisory Group of Experts on Immunization (SAGE), WHO vaccination position papers and technical guidelines, and TAG reports. To date, 7 countries have report having a complete vaccination strategy for mpox – Brazil, Chile, Dominican Republic, Ecuador, El Salvador, Panama, and Uruguay. The following figure describes the number of people who have completed the mpox vaccination schedule in each country that has initiated a vaccination campaign.


**Figure 2.** Number of people who have completed the full mpox vaccination course in each country. Americas Region, as of 18 November 2024



Complete schedule refers to people that have completed the full vaccination course as determined by the country's vaccination policy.  
 Source: PAHO Mpox form updated on 2024/11/18

In addition, demographic information on recipients of mpox vaccination reveals that the major proportion of doses have been received by adults between 19-49 years of age, which corresponds with the age distribution of confirmed cases in the region.

**Figure 3.** Age distribution of people who have completed the full mpox vaccination course. Americas Region, as of 18 November 2024



Complete schedule refers to people that have completed the full vaccination course as determined by the country's vaccination policy.  
Source: PAHO Mpox form updated on 2024/11/18

It is important to take into consideration that, as reported by the Revolving Fund and the WHO, there is limited availability of vaccines and that the vaccines currently available through the RF are already allocated. Given that in the short and medium term, vaccine availability is expected to be very limited, countries are recommended to consider vaccine deployment in phases in their vaccination plans, according to the epidemiological scenario and prioritization of groups at higher risk of severe disease. To this end, it is important to maintain an updated analysis of the Mpox situation in order to guide prevention and control actions, in which vaccination is one of the components.

### Risk Communication and Community Engagement

PAHO has held webinars together with Ministries of Health and organized Civil Societies on topics including Mpox epidemiology, clinical presentations, infection prevention and control, prevention, and treatment.

PAHO has worked with non-governmental organizations, academic institutions, and community-led services working with gay, bisexual, and other men who have sex with men as partners for engagement and risk communication activities with these vulnerable populations. The organization has issued public health recommendations for gay, bisexual, and other men who have sex with men (available on the PAHO website).



The organization has developed and distributed brochures/pamphlets to be used in print and digital with information and general recommendations for the community of gay, bisexual men, and other men who have sex with men to share/distribute with organizers or attendees of festivals and other massive events, and on social media. Flyers with Mpox facts and measures for recovering at home and key information for sex workers were also distributed at healthcare facilities and organizations serving high-risk groups.

PAHO has been monitoring travel measures for mpox through a methodical search across 35 countries in the Region of the Americas. To date, there are no travel measures in any of these countries, which aligns with WHO's recommendations.

Additionally, PAHO has constructed a calendar that categorizes events by type (cultural, sporting, religious, political, and pride) and country. In the first half of November, 3 cultural events (Mexico, Dominican Republic, and Panama), 2 political events (Colombia, and the United States of America), and 7 sporting events (Dominican Republic, Brazil, Venezuela (Bolivarian Republic of), Paraguay, Ecuador, Uruguay, Peru) were observed. For the second half of November, 3 cultural events (Guatemala, Mexico, Panama), 8 sporting events (Bolivia (Plurinational State of), Colombia, Argentina, Chile, Brazil, the United States of America, Paraguay, Peru), and a Pride Parade in Brazil are scheduled.

The WHO has also released two documents: "[Considerations for border health and points of entry for mpox](#)" and "[Gatherings in the context of the 2024 mpox outbreak: Public Health guidance](#)." These documents provide comprehensive advice for managing mpox in these environments, emphasizing coordination, surveillance, and non-discriminatory practices. The first document targets national and subnational health authorities, PoE authorities, public health professionals, civil society organizations, and regional authorities. The second document is aimed at host governments, health authorities, event organizers, healthcare providers, and attendees of meetings of any size and type. These documents are being translated into Spanish and will be sent to the countries, along with the calendar.