

2008 Immunization Schedule for Selected Vaccines - Latin American Countries

	BCG		OPV							DPT					DPT-Hep B+Hib				DPT+Hib				Hep B**				Hib				
	Number of doses																														
Country	1	2	NB§	1	2	3	4	5	6	1	2	3	4	5	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	
ARG	NB			2m	4m	6m	18m	6y						6y	May 2009				2m	4m	6m	18m	NB	2m	6m						
BOL*	NB			2m	4m	6m									2m	4m	6m														
BRA*	NB			2m	4m	6m	15m						15m	4-6y					2m	4m	6m		NB	1m	6m						
CHI	NB			2m	4m	6m	18m						18m	4y	2m	4m	6m														
COL	NB		NB	2m	4m	6m	18m	5y					18m	5y	2m	4m	6m						NB								
COR	NB			2m	4m	6m	4y				4m	15m	4y		2m	6m							NB						4m	15m	
CUB	NB			2 (<1y)	2 (1y)	2 (2y)	1 (9y)						18m	6y*	2m	4m	6m						NB								18m
DOR	NB			2m	4m	6m	18m	4y					18m	4y	2m	4m	6m						NB								
ECU	NB			2m	4m	6m	18m						18m	6y*	2m	4m	6m						NB*	school children at-risk							
ELS	NB			2m	4m	6m	15m	4y					15m	4y	2m	4m	6m						risk groups								
GUT	NB			2m	4m	6m	18m	4y					18m	4y	2m	4m	6m														
HAI	NB		NB	6w	10w	14w	+1y			6w	10w	14w	+1y																		
HON	NB			2m	4m	6m	1-4y						18m	4-5y	2m	4m	6m						NB	risk groups							
MEX*	NB			*	*	*							2y	4y	2m*	4m*	6m*						NB	2m	6m	12y+					
NIC	NB			2m	4m	6m	*						18m		2m	4m	6m						health workers								
PAN	NB			2m	4m	6m	18m	4-5y						4-5y	2m	4m	6m					18m	NB	health workers at risk; school children at risk							
PAR	NB			2m	4m	6m	18m	4y					18m	4y	2m	4m	6m						risk groups								
PER	NB			2m	4m	6m							18m	4y	2m	4m	6m						NB	population at risk							
URU	NB			2m	4m	6m	12m							5y	2m	4m	6m	15m					12y	+1m	+6m						
VEN	NB		NB	2m	4m	6m							18m		2m	4m	6m						NB	risk groups							

text bold and in red are still pending replies

NB: Newborn

WCBA(s): Women of Childbearing Age

** This table only includes the hepatitis B vaccine used in children national schedules. Vaccination of at-risk groups is not included.

*** TT/dT use reported as indicated by countries in the EPI Indicator Tables.

§ Doses administered to newborns do not count towards the completion of the primary series.

* BOL: Hepatitis B for newborns only in selected hospitals.

* CUB: DT pediatric administered at 6 years; vaccine used is tetravalent DPT+Hep B (starting in 2005).

* ECU: DT-2nd booster given to 2nd and 7th graders; HepB given to newborns in municipalities at risk; DT pediatric administered at 6 years

* HON OPV NB dose removed from schedule in 2006

* MEX: OPV doses for children 6 months to 5 years old during National Health Weeks; Pentacel (DTaP-Hib-IPV) given

* NIC: Additional OPV doses for children during National Health Weeks.



2008 Immunization Schedule for Selected Vaccines - Latin American Countries

	M	MMR			MR	TT/dT***					Yellow Fever†	Influenza		Varicella	Hepatitis A	
	Number of Doses															
Country	1	1	2	3	1	1	2	3	4	5	1	1	2	1	1	2
ARG		1y	6y		MEF	16y	every 10y				≥1y in population in border areas	risk groups; ≥65y		risk groups starting 2009	12m	
BOL*		12-23m				WCBAs	WCBAs	WCBAs	WCBAs	WCBAs	12-23m					
BRA*		1y	4-6y		WCBAs	>7y	WCBAs				9m; 10y	≥60y; risk groups		indigenous populations	special groups	
CHI		1y	6y			7y						risk groups; ≥60y; pregnant women; health workers				
COL		1y	5y			WCBAs				1y; population in at-risk areas;	6m-23m (2 doses); ≥65y in some districts					
COR		15m	7y			10y	every 10y					risk groups; 6m to <8y with risk factors; ≥65y		risk groups; 15m		
CUB		1y	6y			15y*						≥65y; children <24m w/chronic diseases				
DOR		1y				WCBAs	WCBAs	WCBAs				>65y; persons w/chronic diseases				
ECU		12-23m			6y	11y	WCBAs				>1y at risk areas	6-23m ; ≥65y				
ELS		1y	4y			≥10y	every 10y pregnant women					risk groups; 6m-23m; >60y				
GUT		1y					WCBAs					>60y				
HAI					9m	WCBAs										
HON		1y			susceptibles	11-12y	WCBAs					6-23m with chronic diseases; risk groups; >60y				
MEX*		1y	6y			11-12y (Grade 6)	WCBAs	WCBAs	WCBAs	WCBAs		6m-35m; ≥60y and risk groups		immunocompromised 1y (social security)		
NIC		1y				6-9y	10-14y	MEF				6-23m; >65y with chronic disease				
PAN		1y	4-5y			6y	MEF				>1y in population at-risk	6m-23m, >60y; risk groups			12m	18m
PAR		1y	4y		5y-39y	WCBAs	WCBAs	WCBAs	WCBAs	WCBAs	≥1y in population in border areas	6-23m; >60; other risk groups				
PER		1y	4y			WCBAs	WCBAs	WCBAs	WCBAs	WCBAs	1y	risk groups				
URU		1y	5y			12y	every 10y					risk groups; 6-23m; ≥65y		1y	15m	21m
VEN		1y				WCBAs	WCBAs	WCBAs	WCBAs	WCBAs	>1y*	≥65y; risk groups				

NB: Newborn

WCBA(s): Women of Childbearing Age

§ Doses administered to newborns do not count towards the completion of the primary series.

† Yellow Fever vaccine administered to travellers to endemic countries not included

* DOR: Influenza given to >60 year olds in retirement homes

* PAN: Introduced Hepatitis A vaccines in May 2006 and rotavirus vaccine in Feb 2006.

* VEN: Yellow fever vaccine administered to population in at-risk municipalities, in addition to recruits and travelers to endemic areas.

* CUB: TT only



2008 Immunization Schedule for Selected Vaccines - Latin American Countries

Country	Rotavirus		Vitamin A									Pneumococcal Conjugate					Meningitis BC	
	1	2	Number of Doses									1	2	3	4	5	1	2
ARG												risk groups						
BOL*	2m	4m		6-11m	1-4y													
BRA*	2m	4m										2m*	4m*	6m*	booster 12-15m*			
CHI																		
COL	to population at risk April 2008											Heptavalente (April 2008); given to population at risk						
COR												2m	4m	6m	15m			
CUB																	3m	5m
DOR				6m	12m	18m	24m	30m										
ECU	2m	4m		6m	12m	18m	24m	30m	36m									
ELS	2m	4m																
GUT				6m	12m	18m	24m	30m	36m	42m	48m	54m						
HAI				6m-7y														
HON	intro in feb 2009			6m	12m	18m	24m	30m	36m									
MEX*	2m	4m		NB	6-59m							2m	4m	12m				
NIC	2m	4m	6m	6m	1y	2y	3y	4y				2m	4m	6m	(intro in 2009)			
PAN	2m	4m		6m	12m	18m						2m*	4m*	6m*	12-15m*			
PAR																		
PER		4m*	6m*	6m	12m	23m						2m*	4m*	6m*				
URU												2m	4m	12m				
VEN	2m	4m																

NB: Newborn

WCBA(s): Women of Childbearing Age

§ Doses administered to newborns do not count towards the completion of the primary series.

* ARG: Tdap will be administered to health workers starting in 2009; Argentinian hemorrhagic fever for groups at risk

* BRA: Pneumo_conj given to population at risk

* ECU: Rotavirus introduced in October 2007

* MEX: Rotavirus will be extended in the national level; SNS-Semana Nacional de Salud; Pneumo_conj given subnationally, to be scaled up in 2008

* PAN: in population at risk

* PER: At subnational



2008 Immunization Schedule for Selected Vaccines - Latin American Countries

Country	Typhoid Fever			Tdap	Pneumo_Polysaccharide	HPV
	1	2	3	1	1	1
ARG	armed forces starting 2009			11a*	risk groups	11a
BOL*						
BRA*	risk groups				risk groups	
CHI					65y+	
COL						
COR				?		
CUB	10y	13y	16y			
DOR						
ECU					older adults	
ELS						
GUT						
HAI						
HON						
MEX*					≥65y	12y; risk municipalities*
NIC						
PAN				11y		10y (3 doses)
PAR						
PER						3 doses*
URU						
VEN						

* MEX: will be given nationwide in 2009

* PER: at subnational level

